

DOI: 10.34921/amj.2026.2.001

MƏDƏNİN YUXARI HİSSƏSİ XƏRÇƏNGİNİN MƏRHƏLƏLƏRİNİN TƏYİNİNDƏ ULTRASƏS MÜAYİNƏSİNİN İMKANLARI

**R.Ya.Abdullayev¹, İ.V.Kriyanovskaya¹, F.İ.İslamzadə², O.A.Nakoneçnaya¹,
F.İ.Kulikova³, J.M.Xvorostenko³, I.N.Kixtenko³, V.F.Zavizion³, D.V.Şaqan⁴**

¹Xarkov Milli Tibb Universiteti, Xarkov, Ukrayna;

²Azərbaycan Tibb Universiteti, Bakı, Azərbaycan;

³Dnepr Milli Tibb Universiteti, Dnepr, Ukrayna;

⁴Otorinolarinqologiya klinikası, Dnepr, Ukrayna

Məqalədə mədənin yuxarı hissəsi xərçənginin diaqnostikasında sonografiyanın effektivliyini araşdırmaq məqsədilə aparılmış tədqiqat işinin nəticələri əks etdirilmişdir. Mədə xərçəngi diaqnozu histoloji müayinə əsasında təsdiq edilmiş 29 pasiyentin müayinəsinin nəticələri araşdırılmışdır. Onlardan 5 nəfərdə T₂, 16 nəfərdə T₃, 8 nəfərdə T₄ mərhələli mədə xərçəngi olmuşdur; sonografiya və kompüter-tomografiya müayinələrinin nəticələri müqayisə edilmişdir.

KT müayinəsinin nəticələri 27 xəstədə (93,1%) mədə xərçənginə dəqiq diaqnoz qoyulması ilə nəticələnmişdir – xərçəngin T₂ mərhələsi olan 4 nəfərdə (13,8%), T₃ mərhələsində olan 15 nəfərdə (51,3%), T₄ mərhələsində olan 8 nəfərdə (27,6%). Sonoqrafiya isə 23 nəfərdə (79,3%) mədə xərçənginin dəqiq diaqnozunu qoymağa imkan vermişdir ki, onlardan da 3-nəfərdə (10,3%) proses T₂, 14 nəfərdə (48,3%) – T₃ və 6 nəfərdə (20,7%) T₄ mərhələsində olmuşdur. KT və sonografiya müayinələrinin nəticələri arasında statistik etibarlı fərq aşkar edilməmişdir.

5 xəstənin qida borusunun abdominal hissəsində stenoz olduğuna görə qastroskopiya aparmaq mümkün olmamışdır. Onlardan hər birində stenoz həm sonografiya, həm də KT vasitəsilə aşkar edilmişdir. Mədənin yuxarı hissəsinin xərçəngi üçün sonografiyanın həssaslığı 87,5%, spesifikliyi 60,0% və dəqiqliyi 82,8% olmuşdur.

Tədqiqata əsasən belə qənaətə gəlmək olar ki, sonografiya ucuz başa gəlidiyinə və qeyri-invaziv üsul olduğuna görə, həm skrining üçün, həm də mədə xərçənginin mərhələsini təyin etmək üçün əlverişli bir üsul kimi istifadə edilə bilər. Bu, fibroqastroskopiyadan istifadəedilmə imkanlarının məhdud olduğu hallarda xüsusilə əhəmiyyətlidir.

Açar sözlər: sonografiya, yuxarı mədə xərçəngi, kompüter tomoqrafiyası

Keywords: sonography, upper gastric carcinoma, computed tomography

Ключевые слова: сонография, рак верхнего отдела желудка, компьютерная томография

POSSIBILITIES OF ULTRASONOGRAPHY IN DETERMINING THE STAGE OF UPPER GASTRIC CANCER

**R.Ya.Abdullaiev¹, I.V.Krizhanovska¹, F.I.Islamzade², O.A.Nakonechna¹, F.I.Kulikova³,
J.M.Khvorostenko³, I.N.Kikhtenko³, V.F.Zavizion³, D.V.Shahan⁴**

¹Kharkiv Nathional Medical University, Ukraine; ²Azerbaijan Medical University, Baku, Azerbaijan; ³Dnipro State Medical University, Dnipro, Ukraine;

⁴LLC Otorhinolaryngology clinic, Dnipro, Ukraine

The article presents the possibilities of sonography in diagnostics of upper gastric cancer. Gastric cancer was diagnosed based on histological examination of biopsy material in 29 patients, including 5 with stage T₂, 16 with stage T₃, and 8 with stage T₄ tumors. Sonography and CT results were compared. Accurate diagnosis of stage T gastric cancer using CT was established in 27 (93.1%) patients, including 4 (13.8%) with stage T₂, 15 (51.7%) with stage T₃, and 8 (27.6%) with stage T₄. Sonography allowed accurate diagnosis of gastric cancer stage in 23 (79.3%) patients, including 3 (10.3%) with stage T₂, 14 (48.3%) with stage T₃, and 6 (20.7%) with stage T₄. No statistically significant differences were found between CT and sonography results. Abdominal esophageal stenosis was detected in 5 patients (17.2%),

preventing gastroscopy. In all cases, stenosis was detected by sonography and CT. Sonography's sensitivity for diagnosing upper gastric cancer was 87.5%, specificity 60.0%, and accuracy 82.8%, respectively.

Based on this study, it can be concluded that sonography, as an inexpensive, noninvasive method, can be used not only for screening but also for staging upper gastric cancer, which in some cases limits the use of fibrogastroscopy.

The effectiveness of treatment for patients with stomach cancer depends on the stage of the tumor. For very early, superficial tumors (T1a), endoscopic mucosal resection/submucosal dissection is the preferred procedure, whereas for early-stage cancers that are not amenable to endoscopic resection, surgical resection is the treatment of choice. Total/distal gastrectomy, depending on the tumor location, in conjunction with neoadjuvant chemotherapy, is the standard treatment for locally advanced GC ($\geq T3$, any N or $\geq T2$, N+). For advanced unresectable/metastatic GC (35–40% of cases at the time of initial diagnosis), chemotherapy is still considered the standard treatment. Therefore, it is crucial to correctly determine the stage of disease in order to select the most effective therapeutic pathway, and imaging plays a pivotal role in this regard [2]. Gastric cancer can spread via the lymphatic system to the perigastric ligaments, mesentery, omentum and adjacent and distant organs [1].

In recent decades, diagnostic imaging, particularly cross-sectional techniques including contrast-enhanced computed tomography (CT), magnetic resonance imaging (MRI), and positron emission tomography (PET), has assumed a central role in the diagnosis of various pathologic entities. In current clinical practice, one of the most common goals is to stage cancer patients in order to best assess their treatment needs and guide them toward surgical or medical interventions [2-3].

Recent publications demonstrate the potential of sonography in diagnosing gastric cancer and determining its location. In diagnosing gastric outlet cancer, the sensitivity and accuracy of ultrasonography is comparable to CT [4-5]. Localization of gastric cardia cancer in the late stages leads to stenosis of the abdominal esophagus, which limits the capabilities of fibrogastroscopy. All this increases the relevance of using sonography for diagnosing gastric cardia and fundus cancer.

Objective: to evaluate the capabilities of sonography in the diagnosis of upper gastric cancer.

Materials and methods: The study is based on observations of 29 patients with gastric cardia and fundus cancer, aged 43 to 74 years (mean age 56 ± 5 years). Of the patients examined, stage T2 gastric cancer was diagnosed in 5 (17.2%), stage T3 in 16 (55.7%), and stage T4 in 8 (27.6%) patients, respectively. All patients underwent radiographic examination, fibrogastroscopy (FGS) with biopsy, multidetector computed tomography (MDCT), and gastric ultrasonography. Ultrasound was performed using convex transducer in the 2-5 MHz frequency range on a Philips HD-11 scanner, and computed tomography was performed on a Toshiba Aquilion 64 system.

To assess differences between quantitative indicators in the compared groups, the Student's t-test and the χ^2 test were used. Differences were considered statistically significant at $p < 0.05$. The sensitivity, specificity, and accuracy of sonography in the diagnosis of upper gastric cancer were studied.

Results. Table 1 shows the distribution of upper gastric cancer cases depending on the T stage. As can be seen from the table, upper gastric cancer at stage T2 was diagnosed in 5 ($17.2 \pm 7.0\%$) patients, at stage T3 – in 16 ($55.2 \pm 9.2\%$) and at stage T4 – in 8 ($27.6 \pm 8.3\%$) patients, respectively. Gastric cancer at stage T3 was diagnosed statistically significantly more often than at stage T2 ($P < 0.01$) and at stage T4 ($P < 0.05$).

Table 1. Distribution of cases of upper gastric cancer taking into account the T stage of the tumor

Cancer stage			
T1	T2	T3	T4
-	n=5	n=16	n=8
-	17,2 $\pm$ 7,0%	55,2 $\pm$ 9,2%	27,6 $\pm$ 8,3%
-		P<0,05 P<0,01	
	n=29		

Since computed tomography is currently the main method for determining the stage and extent of gastric cancer, we compared their results in order to evaluate the diagnostic capability of sonography (Table 2). When determining the stage of gastric cancer based on surgical data, the results of CT coincided in 27 ($93.1 \pm 4.7\%$) cases, and USG – in 23 ($79.3 \pm 7.5\%$) cases. At stage T2, gastric carci-

Table 2. Comparison of ultrasonography and CT results taking into account the stage of gastric cancer


Methods	Upper gastric cancer stage (n = 29)		
	T2 (n=5)	T3 (n=16)	T4 (n=8)
KT	4 (13,8±13,2%)	15 (51,7±9,3%)	8 (27,6±8,3%)
YCF	3 (10,3±5,6%)	14 (48,3±9,3%)	6 (20,7±7,5%)

noma was detected by CT in 4 (13.8±13.2%) cases, and by sonography – in 3 (10.3±5.6%) cases, at stage T3 – in 15 (51.7±9.3%) and 14 (48.3±9.3%) cases, at stage T4 – in 8 (27.6±8.3%) and 6 (20.7±7.5%) cases, respectively.

Table 3 presents the diagnostic performance of sonography for gastric cardia and fundus cancer. In diagnosing gastric cancer, sonography sensitivity was 87.5%, specificity 60.0%, accuracy 82.8%, positive predictive value 91.3%, and negative predictive value 50.0%.

Due to the specific direction of ultrasound waves to the object being examined, the cardiac region of the stomach, even with minimal tumor infiltration, was always visualized more clearly than the fundus. When tumor spread to the abdominal esophagus due to the development of stenosis, fibrogastroscopy was difficult and was not performed in 5 cases. In diffusely infiltrative gastric cancers with a smooth mucosal surface, the main tumor feature was thickening of the muscular layer with preserved differentiation between the serous

and mucosal membranes. When stenosis of the abdominal esophagus develops, the narrowed lumen is better defined during fluid passage. Figures 1 and 2 show sonographic and CT imaging of cancer of the cardiac region of the stomach.

**Fig. 1.** Sonographic visualization of gastric cardia cancer with extension into the esophagus (vertical arrows). The horizontal arrow indicates the narrowed lumen of the abdominal esophagus.**Table 3.** CT and Sonography results in the diagnosis of upper gastric cancer

CT (n=29)				USG (n=29)			
TP	FN	TN	FP	TP	FN	TN	FP
24	2	2	1	21	3	3	2
Sensitivity		92,3%		Sensitivity		87,5%	
Specificity		66,7%		Specificity		60,0%	
Accuracy		89,7%		Accuracy		82,8%	
Positive predictive value		96,0%		Positive predictive value		91,3%	
Negative predictive value		50,0%		Negative predictive value		50,0%	

Note: TP – true positive, FN – false negative, TN – true negative, FP – false positive




Fig. 2. CT visualization of gastric upper cancer.




Fig. 3. Gastric fundus cancer at stage T2. Arrows indicate localized thickening of the gastric wall. Ultrasound imaging of the tumor was only possible after a targeted search for it following diagnostic testing with fibrogastroscopy and CT.

Examination of the fundus was best accomplished in the Trendelenburg position. At all stages of fundus cancer, the wall layers were undifferentiated. At stage T2, only gastric cancer already diagnosed during fibrogastroscopy could be visualized using ultrasonography (Fig. 3).

Discussion

The absence of clinical symptoms in the early stages of gastric cancer makes its timely diagnosis difficult [6]. It is known that radiography and endoscopy are the most common methods for diagnosing this pathology [7-8]. At the same time, in the last two decades, gastric cancer staging has been carried out using computed tomography. The method has several advantages over other imaging methods in detecting tumor invasion beyond the wall of these organs and determining metastases to the lymph nodes. According to some experts, even CT does not always allow differentiating between T2/T3 stages of gastric cancer. It is preferable to use various imaging methods, each of which may have several advantages over the others in each specific case [9].

A recently published meta-analysis highlights the good diagnostic value for the

detection of peritoneal carcinomatosis, with pooled sensitivity and specificity values of 89% and 86%, respectively. When included studies were grouped by primary tumor, a pooled sensitivity of 97% was reported for gastrointestinal malignancies [10].

Finally, a diagnostic meta-analysis was used to determine the accuracy of CT and endoscopic ultrasound in the staging of gastric cancer. The results indicated that endoscopic ultrasound is superior to CT for T1 staging, whereas no significant differences were found for T2–T4 lesions or stage N1 [11].

In recent years, advances in ultrasound diagnostic technology have enabled the combined use of grayscale, Doppler, and elastographic modes, significantly expanding the capabilities of the method and, in some cases, offering advantages over CT and MRI [18]. Previously published studies have shown that the results of ultrasonography and CT in diagnosing gastric cancer do not significantly differ [12]. In late-stage gastric and colon cancer, stenosis of the lumen of these hollow organs prevents endoscopy [13]. In such cases, the capabilities of ultrasound examination are significantly expanded, offering an accessible, non-invasive, inexpensive, and

informative method for diagnosing gastric cancer. As a result of our research, the sensitivity, specificity, and accuracy of transabdominal ultrasonography have approached those of CT.

Conclusions

1. For upper gastric cancer, the diagnostic performance of ultrasonography is only

slightly inferior to that of computed tomography, while its non-invasiveness and low cost allow it to be used for tumor staging.

2. Considering that abdominal esophageal stenosis develops in late-stage upper gastric cancer, which limits the use of fibrogastroscopy, the expanded use of ultrasonography is encouraged.

Conflict of interest

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

REFERENCES

1. Young J.J., Pahwa A., Patel M. et al. Ligaments and Lymphatic Pathways in Gastric Adenocarcinoma // RadioGraphics. 2019, №39, p.668-689. doi: 10.1148/rg.2019180113.
2. Saiz Martínez R., Dromain C., Vietti Violi N. Imaging of Gastric Carcinomatosis // J. Clin. Med. 2021, №10, p.5294. doi: 10.3390/jcm10225294.
3. Ajani J.A., D'Amico T.A., Bentrem D.J. et al. Gastric Cancer, Version 2.2022, NCCN Clinical Practice Guidelines in Oncology // J. Natl. Compr. Canc. Netw. 2022, №20, 167-192. doi: 10.6004/jnccn.2022.0008.
4. Nie R.C., Yuan S.Q., Chen X.J. et al. Endoscopic ultrasonography compared with multidetector computed tomography for the preoperative staging of gastric cancer: a meta-analysis // World J. Surg. Oncol. 2017, Vol. 15, p. 113. doi: 10.1186/s12957-017-1176-6.
5. Aliev C.A., Abdullaev R.Ya., Kryzhanovskaya I.V. The role of ultrasonography in diagnosis of gastric cancer localization and stage // Azerbaijan Medical Journal, 2019, №3, 51-59.
6. Sung H., Ferlay J., Siegel R.L., et al. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries // CA Cancer J Clin. 2021, Vol.71, №3, p.209-249. doi: 10.3322/caac.21660.
7. Yashima K., Shabana M., Kurumi H. et al. Gastric cancer screening in Japan: a narrative review // J Clin Med. 2022;11(15):4337. doi: 10.3390/jcm11154337. Published 2022 Jul 26.
8. Mabe K., Inoue K, Kamada T, Kato K, Kato M, Haruma K. Endoscopic screening for gastric cancer in Japan: current status and future perspectives. Dig Endosc. 2022;34(3):412-419. doi: 10.1111/den.14063.
9. Sharara S.M., Nagi M.A., Soliman S.S. (2018) Multidetector computed tomography in the evaluation of gastric malignancy; a multicenteric study. The Egyptian Journal of Radiology and Nuclear Medicine 49(2):304–309
10. Dong L., Li K., Peng T. Diagnostic Value of Diffusion-Weighted Imaging/Magnetic Resonance Imaging for Peritoneal Metastasis from Malignant Tumor: A Systematic Review and Meta-Analysis. Medicine. 2021;100:e24251. doi: 10.1097/MD.00000000000024251.
11. Fan Z., Cong Y., Zhang Z., Li R., Wang S., Yan K. Shear Wave Elastography in Rectal Cancer Staging, Compared with Endorectal Ultrasonography and Magnetic Resonance Imaging. Ultrasound Med. Biol. 2019, 45, 1586-1593.
12. Abdullaev R.Ya., Dikan İ.N., Babkina T.M., Kulikova F.İ., Qolovko T.S., Solodyanikova O.İ., Dudnik T.A. Ultrasonography in the examination of patients with gastric cancer: Comparison with MDCT results. Azerbaijan Medical Journal, 2020, №3, 12-19.
13. Abdullaev R.Ya., Kulikova F.J., Golovko T.S. et al. Possibilities of transabdominal ultrasonography in the diagnosis of gastric and colon cancer // Azerbaijan Medical Journal, 2023, №1, p.10-16. <https://doi.org/10.34921/atj.2023.1.001>

ВОЗМОЖНОСТИ УЛЬТРАСОНОГРАФИИ В ОПРЕДЕЛЕНИИ СТАДИЙ РАКА ВЕРХНЕГО ОТДЕЛА ЖЕЛУДКА

Р.Я.Абдуллаев¹, И.В.Крыжановская¹, Ф.И.Исламзаде², О.А.Наконечная¹, Ф.И.Куликова³,
Ю.М.Хворостенко³, И.Н.Кихтенко³, В.Ф.Завизион³, Д.В. Шаган⁴

¹Харьковский национальный медицинский университет, Харьков, Украина;

²Азербайджанский медицинский университет, Баку, Азербайджан;

³Днепропетровский государственный медицинский университет, Днепр, Украина;

⁴ТОВ «ЛОП-Клиника», Днепр, Украина

Резюме. В статье показаны возможности сонографии в диагностике рака верхнего отдела желудка. Рак желудка был установлен на основании гистологического исследования биопсийного материала у 29 пациентов, у 5 из них опухоль была на стадии Т2, у 16 – на стадии Т3 и у 8 – на стадии Т4. Были сопоставлены результаты сонографии и КТ. Точная диагностика Т стадии рака желудка с помощью КТ была установлена у 27 (93,1%) больных – из них у 4 (13,8%) на стадии Т2, у 15 (51,7%) на стадии Т3 и у 8 (27,6%) на стадии Т4. Сонография позволила точно диагностировать стадию рака желудка у 23 (79,3%) больных – у 3 (10,3%) на стадии Т2, у 14 (48,3%) на стадии Т3 и у 6 (20,7%) на стадии Т4. Статистически достоверных различий между результатами КТ и сонографии не было выявлено. У 5 (17,2%) пациентов был выявлен стеноз абдоминального отдела пищевода, из-за которого не проводилась гастрфиброскопия. Во всех случаях стеноз был выявлен при сонографии и КТ. В диагностике рака верхнего отдела желудка чувствительность сонографии составила 87,5%, специфичность 60,0%, точность 82,8%, соответственно.

На основании проведенного исследования можно сделать вывод о том, что сонография являясь недорогим, неинвазивным методом может быть использована не только для скрининга, а также для определения стадии верхнего отдела рака желудка, при которых в некоторых случаях ограничивается применение фиброгастроскопии.

For correspondence:

Abdullaiev Rizvan Yagubovich, Department of Ultrasound of Kharkiv National Medical University, Ukraine

E-mail: rizvanabdullaiev@gmail.com

Тəqdim edilmişdir: 21.01.2026

Qəbul edilmişdir: 14.04.2026